

Molecular outflow in the composite galaxy NGC 6764

Stéphane Leon

*Instituto de Astrofísica de Andalucía, Camino Bajo de Huétor, 24 Apdo
 3004, 18080 Granada, Spain*

Andreas Eckart

I. Physikalisches Institut, University of Cologne, Cologne, Germany

Seppo Laine

*SIRTF Science Center, California Institute of Technology, 220-6,
 Pasadena, CA 91125*

Eva Schinnerer

*Department of Astronomy, California Institute of Technology,
 Pasadena, CA 91125-2400, USA*

Abstract. NGC 6764 is a composite galaxy exhibiting central activity (LINER) and recent massive star formation (Wolf-Rayet galaxy). We report new high-resolution interferometric data of the $^{12}\text{CO}(1-0)$ and $^{12}\text{CO}(2-1)$ emission in the central part of the galaxy together with high resolution VLA data of the radio continuum emission. The bulk of molecular gas (few times $10^8 M_{\odot}$) is located in the central part. A molecular outflow ($\sim 1.7 \times 10^7 M_{\odot}$) out-of the plane is associated with the starburst-driven outflow radiocontinuum emission.

1. Introduction

NGC 6764 is a barred spiral galaxy classified as a low-ionization nuclear emission-line region (LINER) from optical spectroscopy. This nearby active galaxy ($D=32$ Mpc for $H_0 = 75 \text{ Mpc}^{-1} \text{ km s}^{-1}$; $1'' = 160 \text{ pc}$) is interesting as well for its nuclear starburst which is few Myr old as shown by its Wolf-Rayet features at $0.47 \mu\text{m}$ (HeII). This galaxy has been studied in details in previous papers (Eckart et al. 1991,1996; Schinnerer et al. 2000) at different wavelengths (NIR, X-ray, single dish CO). The study of NGC 6764 has been motivated by the composite nature of its nucleus hosting a young powerful starburst as well as an AGN.

2. Results

The $^{12}\text{CO}(1-0)$ and $^{12}\text{CO}(2-1)$ lines have been observed with the IRAM interferometer located on the Plateau de Bure (France) with a spatial resolution of $2.0'' \times 1.6''$ (PA= 7°) for the $^{12}\text{CO}(1-0)$ data and $1.25'' \times 1.25''$ for the $^{12}\text{CO}(2-1)$




Figure 1. VLA 20cm emission overlaid by $^{12}\text{CO}(1-0)$ emission with contour levels from 1 to 27 $\text{Jy.beam}^{-1}\text{km/s}$ by step of 1 $\text{Jy.beam}^{-1}\text{km/s}$. The position of the major axis and the bar are shown respectively by the solid and dashed lines.

data. CO emission has been found in the very center of the galaxy ($R < 1.5$ kpc, about $10''$) and the total mass of molecular gas mass (including He) is found to be about $6.7 \times 10^8 M_{\odot}$. About 50 % of the total molecular gas mass is concentrated in one $^{12}\text{CO}(1-0)$ beamsize in the center of NGC 6764. An offset in the position of the CO peak and the 20cm radio continuum peak is observed ($\sim 1.2''$). The most striking feature is the molecular gas flaring above the plane correlated with the 20cm radio continuum outflow starburst-driven (see Fig. 1). The $^{12}\text{CO}(1-0)$ line is observed up to 1 kpc above the galactic plane. An estimate gives a molecular gas mass of $1.7 \times 10^7 M_{\odot}$ out-of the plane, with $z > 500$ pc (~ 1.5 CO(1-0)-beamsize). The physical conditions of the molecular gas is probed by the CO(2-1)/CO(1-0) line ratio which is found to be about 1–1.3. Nevertheless a peak of about 2.0 for this line ratio is found, spatially associated with the active nucleus, at a blueshifted velocity of -140 km/s (rel. to the galaxy) when a lower CO line ratio is found outside the central part (mainly the bar). The nuclear CO kinematics show some molecular gas associated with bar orbits (x_2 orbits).

References

- Eckart, A., Cameron, M., Jackson, J. M., Genzel, R., Harris, A. I., Wild, W., Zinnecker, H. 1991, ApJ, 372, 67
- Eckart, A., Cameron, M., Boller, Th., Krabbe, A., Blietz, M., Nakai, N., Wagner, S. J., Sternberg, A. 1996, ApJ, 472, 588
- Schinnerer, E., Eckart, A., Boller, Th. 2000, ApJ., 545, 205